

UNIQUE DETERMINATION OF THE DAMPING COEFFICIENT IN THE WAVE EQUATION USING POINT SOURCE AND RECEIVER DATA

MANMOHAN VASHISTH

ABSTRACT. In this article, an inverse problems of determining the damping coefficient appearing in a wave equation is studied. We proved that the time-derivative perturbation of wave operator in three space dimension can be recovered uniquely from the data coming from a single coincident source-receiver pair. Since the problem under consideration is under-determined, so some extra assumption on the coefficient is required to prove the uniqueness.

Keywords : Inverse problems, wave equation, point source-receiver, damping coefficient

Mathematics subject classification 2010: 35L05, 35L10, 35R30, 74J25

1. INTRODUCTION

In this article, we consider an inverse problem related to an initial value problem (IVP) for wave operator given by

$$\begin{aligned}(\square - q(x)\partial_t)u(x, t) &= \delta(x, t), \quad (x, t) \in \mathbb{R}^3 \times \mathbb{R} \\ u(x, t)|_{t < 0} &= 0, \quad x \in \mathbb{R}^3\end{aligned}\tag{1}$$

where $\square := \partial_t^2 - \Delta_x$ denotes the wave operator and the coefficient $q \in C^\infty(\mathbb{R}^3)$ is known as damping coefficient. In this paper, we study the problem of determination of coefficient q appearing in (1) from the knowledge of solution measured at a single point for a certain period of time. We are interested in the uniqueness of determination of coefficients q from the knowledge of $u(0, t)$ for $t \in [0, T]$ with $T > 0$ in Equation (1). The problem studied here is motivated by geophysics, where geophysicists wish to determine the properties of earth structure by sending the waves from the surface of the earth and measuring the corresponding scattered responses (see [2, 24] and references therein). Since the coefficient to be determined here depends on three variables while the given data depends on one variable as far as the parameter count is concerned, the problem studied here is under-determined. Thus some extra assumptions on coefficient q are required in order to make the inverse problem solvable. We prove the uniqueness result for the radial coefficient.

There are several results related to the inverse problems for the wave equation with point source. We list them here. Romanov in [18] considered the problem for determining the damping and potential coefficient in the wave equation with point source and proved unique determination of these coefficients by measuring the solution on a set containing infinite points. In [12] the problem of determining the radial potential from the knowledge of solution measured on a unit sphere for some time interval is studied. Rakesh and Sacks in [16] established the uniqueness for angular controlled potential in the wave equation from the knowledge of solution and its radial derivative measured on a unit sphere. In the above mentioned works the measurement set is an infinite set. Next we mention the work where uniqueness is established from the measurement of solution at a single point. Determination of the potential from the data coming from a single coincident source-receiver pair is considered in [15] and the uniqueness result is established for the potentials which are either radial with respect a point different from source location or the potentials which are comparable. Recently author in [25] extended the result of [15] to a separated point source and receiver data. To the best of our understanding, very few results exist in the literature involving the recovery of the damping coefficient from point source and receiver data. Our result, Theorem 1.1, is work in this

direction. In the 1-dimensional inverse problems context, several results exist involving the uniqueness of recovery of the coefficient which depends on the space variable corresponding to the first order derivative; see [9, 10, 11, 13, 19, 22]. We refer to [1, 3, 8, 14, 17] and references therein for more works related to the point source inverse problems for the wave equation.

We now state the main results of this article.

Theorem 1.1. *Suppose $q_i(x) \in C^\infty(\mathbb{R}^3)$, $i = 1, 2$ with $q_i(x) = A_i(|x|)$ for some C^∞ function A_i on $[0, \infty)$. Let u_i be the solution of the IVP*

$$\begin{aligned} (\square - q_i(x)\partial_t)u_i(x, t) &= \delta(x, t) & (x, t) \in \mathbb{R}^3 \times \mathbb{R} \\ u_i(x, t)|_{t < 0} &= 0 & x \in \mathbb{R}^3. \end{aligned} \quad (2)$$

If $u_1(0, t) = u_2(0, t)$ for all $t \in [0, T]$ for some $T > 0$, then $q_1(x) = q_2(x)$ for all x with $|x| \leq T/2$, provided $q_1(0) = q_2(0)$.

The proof of the above theorem is based on an integral identity derived using the solution to an adjoint problem as used in [21] and [23]. This idea was used in [4, 17, 25] as well.

Remark 1.2. Since in Theorem 1.1 we have assumed that the damping coefficient q is radial therefore using the spherical means Equation (2) can be reduced to the one dimensional wave equation. And the one dimensional problem is closely related to the problem studied in [9]. They proved the determination of coefficients by establishing an intrinsic relation between the given data and the inverse spectral theory. In the present paper we are interested in studying the problem in three space dimension where this problem becomes an under-determined problem because the given data depend on one dimension and the coefficient to be determined depend on three dimensions. In this paper, we have provided another proof for determining coefficient q . We believe that our proof is simpler than that of [9] and we do not need to use any boundary constraints for proving Theorem 1.1 as compare to [9].

The article is organized as follows. In Section 2, we state the existence and uniqueness results for the solution of Equation (1), the proof of which is given in [5, 8, 20]. Section 3 contains the proof of Theorem 1.1.

2. PRELIMINARIES

Proposition 2.1. [5, pp.139,140] *Suppose $q \in C^\infty(\mathbb{R}^3)$ and $u(x, t)$ satisfies the following initial value problem*

$$\begin{aligned} Pu(x, t) &:= (\square - q(x)\partial_t)u(x, t) = \delta(x, t), & (x, t) \in \mathbb{R}^3 \times \mathbb{R} \\ u(x, t)|_{t < 0} &= 0, & x \in \mathbb{R}^3 \end{aligned} \quad (3)$$

then $u(x, t)$ is given by

$$u(x, t) = \frac{R(x, t)\delta(t - |x|)}{4\pi|x|} + v(x, t) \quad (4)$$

where $v(x, t) = 0$ for $t < |x|$ and in the region $t > |x|$, $v(x, t)$ is a C^∞ solution of the characteristic boundary value problem (Goursat Problem)

$$\begin{aligned} Pv(x, t) &= 0, \text{ for } t > |x| \\ v(x, |x|) &= -\frac{R(x, |x|)}{8\pi} \int_0^1 \frac{PR(sx, s|x|)}{R(sx, s|x|)} ds, \quad \forall x \in \mathbb{R}^3 \end{aligned} \quad (5)$$

and $R(x, t)$ is given by [5, pp. 134]

$$R(x, t) = \exp \left(-\frac{1}{2} \int_0^1 q(sx) t ds \right). \quad (6)$$

3. PROOF OF THEOREM 1.1

In this section, we prove Theorem 1.1. We will first prove an integral identity which will be used to prove our main result.

Lemma 3.1. *Let $u_i(x, t)$ for $i = 1, 2$ be the solution to Equation (2). Then the following integral identity holds for all $\sigma \geq 0$*

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t u_2(x, t) u_1(x, 2\sigma - t) dt dx = u(0, 2\sigma) \quad (7)$$

where $q(x) := q_1(x) - q_2(x)$ and $u(x, t) = (u_1 - u_2)(x, t)$.

Proof. Here we have u satisfies the following IVP

$$\begin{aligned} \square u(x, t) - q_1(x) \partial_t u(x, t) &= q(x) \partial_t u_2(x, t) & (x, t) \in \mathbb{R}^3 \times \mathbb{R} \\ u(x, t)|_{t < 0} &= 0 & x \in \mathbb{R}^3. \end{aligned} \quad (8)$$

Multiplying Equation (8) by $u_1(x, 2\sigma - t)$ and integrating over $\mathbb{R}^3 \times \mathbb{R}$, we have

$$\begin{aligned} \int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t u_2(x, t) u_1(x, 2\sigma - t) dt dx &= \int_{\mathbb{R}^3} \int_{\mathbb{R}} (\square u(x, t) - q_1(x) \partial_t u(x, t)) u_1(x, 2\sigma - t) dt dx \\ &= \int_{\mathbb{R}^3} \int_{\mathbb{R}} u(x, t) (\square u(x, t) - q_1(x) \partial_t u_1(x, 2\sigma - t)) dx dt \end{aligned}$$

where in the last step above we have used integration by parts and the properties of v in Proposition 2.1. Thus finally using the fact that u_1 is solution to (2), we get

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t u_2(x, t) u_1(x, 2\sigma - t) dt dx = u(0, 2\sigma); \text{ for all } \sigma \geq 0.$$

This completes the proof of the lemma. □

Using Lemma 3.1 and the fact that $u(0, t) = 0$ for all $t \in [0, T]$, we see that

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t u_2(x, t) u_1(x, 2\sigma - t) dt dx = 0; \text{ for all } \sigma \in [0, T/2].$$

Now using Equation (4), we get

$$\begin{aligned} &\int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t \left(\frac{R_2(x, t) \delta(t - |x|)}{4\pi|x|} + v_2(x, t) \right) \left(\frac{R_1(x, 2\sigma - t) \delta(2\sigma - t - |x|)}{4\pi|x|} \right) dt dx \\ &+ \int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) v_1(x, 2\sigma - t) \partial_t \left(\frac{R_2(x, t) \delta(t - |x|)}{4\pi|x|} + v_2(x, t) \right) dt dx = 0. \end{aligned}$$

This gives

$$\begin{aligned}
& \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}} \frac{q(x) \partial_t R_2(x, t) R_1(x, 2\sigma - t) \delta(t - |x|) \delta(2\sigma - t - |x|)}{16\pi^2 |x|^2} dt dx}_{I_1} \\
& + \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}} \frac{q(x) R_2(x, t) R_1(x, 2\sigma - t) \partial_t \delta(t - |x|) \delta(2\sigma - t - |x|)}{16\pi^2 |x|^2} dt dx}_{I_2} \\
& + \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t \left(\frac{R_2(x, t) \delta(t - |x|)}{4\pi |x|} \right) v_1(x, 2\sigma - t) dt dx}_{I_3} \\
& + \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}} \frac{q(x) \partial_t v_2(x, t) R_1(x, 2\sigma - t) \delta(2\sigma - t - |x|)}{4\pi |x|} dt dx}_{I_4} \\
& + \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t v_2(x, t) v_1(x, 2\sigma - t) dt dx}_{I_5} = 0; \text{ for all } \sigma \in [0, T/2].
\end{aligned} \tag{9}$$

In a compact form, this can be written as

$$I_1 + I_2 + I_3 + I_4 + I_5 = 0. \tag{10}$$

Next we simplify each I_j with $j = 1, 2, \dots, 5$. We will use the fact that $v_i(x, t) = 0$ for $t < |x|$.

We have

$$\begin{aligned}
I_1 &= \int_{\mathbb{R}^3} \int_{\mathbb{R}} \frac{q(x) \partial_t R_2(x, t) R_1(x, 2\sigma - t) \delta(t - |x|) \delta(2\sigma - t - |x|)}{16\pi^2 |x|^2} dt dx \\
&= \int_{|x|=\sigma} \frac{q(x) \partial_t R_2(x, |x|) R_1(x, |x|)}{16\pi^2 |x|^2} dS_x \\
&= - \int_{|x|=\sigma} \frac{q(x) R_1(x, |x|) R_2(x, |x|)}{32\pi^2 |x|^2} \left(\int_0^1 q_2(sx) ds \right) dS_x.
\end{aligned}$$

Next we simplify the integral I_2 . We use the following formula [7, Page 231, Eq.(10)]

$$\int \delta'(r - |x|) \varphi dx = \frac{-1}{|x|^2} \int_{|x|=r} \frac{\partial}{\partial r} (\varphi r^2) dS_x. \tag{11}$$

Note that from this formula, by a change of variable, we have

$$\int \delta'(2r - 2|x|) \varphi dx = \frac{-1}{2|x|^2} \int_{|x|=r} \frac{\partial}{\partial r} (\varphi r^2) dS_x. \tag{12}$$

Now

$$\begin{aligned}
I_2 &= \int_{\mathbb{R}^3} \int_{\mathbb{R}} \frac{q(x)R_2(x, t)R_1(x, 2\sigma - t)\partial_t\delta(t - |x|)\delta(2\sigma - t - |x|)}{16\pi^2|x|^2} dt dx \\
&= \int_{\mathbb{R}^3} \int_{\mathbb{R}} \frac{q(x)R_2(x, t)R_1(x, 2\sigma - t)\delta'(t - |x|)\delta(2\sigma - t - |x|)}{16\pi^2|x|^2} dt dx \\
&= \int_{\mathbb{R}^3} \frac{q(x)R_2(x, 2\sigma - |x|)R_1(x, |x|)\delta'(2\sigma - 2|x|)}{16\pi^2|x|^2} dx \\
&= -\frac{1}{32\pi^2\sigma^2} \int_{|x|=\sigma} \frac{\partial}{\partial r} \{q(x)R_1(x, |x|)R_2(x, 2\sigma - |x|)\} dS_x.
\end{aligned}$$

In the last step above, we used Equation (12).

Next we have

$$I_3 = \int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x)\partial_t \left(\frac{R_2(x, t)\delta(t - |x|)}{4\pi|x|} \right) v_1(x, 2\sigma - t) dx dt.$$

We can view the derivative above as a limit of the difference quotients in the distribution topology [6, pp.48]. Combining this with the fact that v_1 is C^2 in $\{(x, t) : |x| \leq t\}$, we get,

$$\begin{aligned}
I_3 &= - \int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \frac{R_2(x, t)\delta(t - |x|)}{4\pi|x|} \partial_t \left(v_1(x, 2\sigma - t) \right) dx dt \\
&= \int_{\mathbb{R}^3} \frac{q(x)R_2(x, |x|)\partial_t v_1(x, 2\sigma - |x|)}{4\pi|x|} dx.
\end{aligned}$$

Again using the fact that $v_1(x, t) = 0$ for $t < |x|$, we get,

$$I_3 = \int_{|x| \leq \sigma} \frac{q(x)R_2(x, |x|)\partial_t v_1(x, 2\sigma - |x|)}{4\pi|x|} dx.$$

Next we simplify I_4 . Similar to I_3 , we have

$$\begin{aligned}
I_4 &= \int_{\mathbb{R}^3} \int_{\mathbb{R}} \frac{q(x)\partial_t v_2(x, t)R_1(x, 2\sigma - t)\delta(2\sigma - t - |x|)}{4\pi|x|} dt dx \\
&= \int_{|x| \leq \sigma} \frac{q(x)R_1(x, |x|)\partial_t v_2(x, 2\sigma - |x|)}{4\pi|x|} dx.
\end{aligned}$$

Finally, we have

$$\begin{aligned} I_5 &= \int_{\mathbb{R}^3} \int_{\mathbb{R}} q(x) \partial_t v_2(x, t) v_1(x, 2\sigma - t) dt dx \\ &= \int_{|x| \leq \sigma} \int_{|x|}^{2\sigma - |x|} q(x) \partial_t v_2(x, t) v_1(x, 2\sigma - t) dt dx. \end{aligned}$$

Now, we use the fact that q_i is a radial function, that is, $q_i(x) = A_i(|x|)$. Then note that

$$R_i(x, |x|) = \exp\left(-\frac{|x|}{2} \int_0^1 q_i(s|x|) ds\right) = \exp\left(-\frac{|x|}{2} \int_0^1 A_i(s|x|) ds\right)$$

is also radial. For simplicity, we denote $R(x, |x|)$ by $R(|x|)$.

With this, we have

$$I_1 = -\frac{A(\sigma)R_1(\sigma)R_2(\sigma)}{8\pi} \int_0^1 A_2(s\sigma) ds.$$

Next we consider I_2 . First let us consider the derivative:

$$D_r := \frac{\partial}{\partial r} (A(r)R_1(x, r)R_2(x, 2\sigma - r)).$$

After a routine calculation, we get,

$$\begin{aligned} D_r &= A'(r)R_1(x, r)R_2(x, r) - \frac{1}{2}A(r)^2R_1(x, r)R_2(x, 2\sigma - r) \\ &\quad - \sigma A(r)R_1(x, r)R_2(x, 2\sigma - r) \int_0^1 A_2'(rs) ds \\ &= A'(r)R_1(x, r)R_2(x, r) - \frac{1}{2}A(r)^2R_1(x, r)R_2(x, 2\sigma - r) \\ &\quad - A(r)R_1(x, r)R_2(x, 2\sigma - r) \left[\frac{\sigma}{r} \left(A_2(r) - \int_0^1 A_2(rs) ds \right) \right]. \end{aligned}$$

On $|x| = \sigma$, we have

$$\begin{aligned} D_r|_{|x|=\sigma} &= R_1(\sigma)R_2(\sigma) \left[A'(\sigma) - \frac{1}{2}A(\sigma)^2 - A(\sigma)A_2(\sigma) + A(\sigma) \int_0^1 A_2(s\sigma) ds \right] \\ &= R_1(\sigma)R_2(\sigma) \left[A'(\sigma) - \frac{1}{2}A(\sigma)(A_1 + A_2)(\sigma) + A(\sigma) \int_0^1 A_2(s\sigma) ds \right]. \end{aligned}$$

Hence

$$I_2 = -\frac{1}{8\pi} \left(R_1(\sigma)R_2(\sigma) \left[A'(\sigma) - \frac{1}{2}A(\sigma)(A_1 + A_2)(\sigma) + A(\sigma) \int_0^1 A_2(s\sigma) ds \right] \right).$$

Let us denote

$$\tilde{A}(\sigma) = A(\sigma)R_1(\sigma)R_2(\sigma).$$

Then

$$I_2 = -\frac{1}{8\pi} \frac{d}{d\sigma} \tilde{A}(\sigma) - \frac{1}{8\pi} \tilde{A}(\sigma) \int_0^1 A_2(s\sigma) ds.$$

Therefore

$$I_1 + I_2 = -\frac{1}{8\pi} \left(2\tilde{A}(\sigma) \int_0^1 A_2(s\sigma) ds + \frac{d}{d\sigma} \tilde{A}(\sigma) \right).$$

Considering the following integrating factor for $I_1 + I_2$

$$\exp \left(2 \int_0^\sigma \int_0^1 A_2(ts) dt ds \right),$$

we have

$$I_1 + I_2 = -\frac{1}{8\pi} \exp \left(-2 \int_0^\sigma \int_0^1 A_2(ts) dt ds \right) \frac{d}{d\sigma} \left[\exp \left(2 \int_0^\sigma \int_0^1 A_2(ts) dt ds \right) \tilde{A}(\sigma) \right].$$

Now from Equation (10), we have

$$\begin{aligned} & \frac{1}{8\pi} \frac{d}{d\sigma} \left[\tilde{A}(\sigma) \exp \left(2 \int_0^\sigma \int_0^1 A_2(st) ds dt \right) \right] \\ &= \exp \left(2 \int_0^\sigma \int_0^1 A_2(st) ds dt \right) \left[\int_{|x| \leq \sigma} \frac{q(x)R_2(x, |x|)\partial_t \{R_1 v_1\}(x, 2\sigma - |x|)}{4\pi|x|} dx \right. \\ & \quad + \int_{|x| \leq \sigma} \frac{q(x)R_1(x, |x|)\partial_t v_2(x, 2\sigma - |x|)}{4\pi|x|} dx \\ & \quad \left. + \int_{|x| \leq \sigma} \int_{|x|}^{2\sigma - |x|} q(x)\partial_t v_2(x, t)v_1(x, 2\sigma - t) dt dx \right] \text{ for all } \sigma \in [0, T/2]. \end{aligned} \tag{13}$$

Integrating on both sides with respect to σ under the assumption that $\tilde{A}(0) = 0$, we get

$$\begin{aligned} & \exp\left(\int_0^{\tilde{\sigma}} \int_0^1 2A_2(st) ds dt\right) \tilde{A}(\tilde{\sigma}) \\ &= \int_0^{\tilde{\sigma}} \exp\left(\int_0^{\sigma} \int_0^1 2A_2(st) ds dt\right) \left\{ \int_{|x|\leq\sigma} \frac{q(x)R_2(x, |x|)\partial_t v_1(x, 2\sigma - |x|)}{4\pi|x|} dx \right. \\ &+ \int_{|x|\leq\sigma} \frac{q(x)R_1(x, |x|)\partial_t v_2(x, 2\sigma - |x|)}{4\pi|x|} dx \\ &+ \left. \int_{|x|\leq\sigma} \int_{|x|}^{2\sigma-|x|} q(x)\partial_t v_2(x, t)v_1(x, 2\sigma - t) dt dx \right\} d\sigma, \quad \text{for all } \tilde{\sigma} \in [0, T/2]. \end{aligned}$$

Now using the fact that R'_i 's are continuous, non-zero functions, and v'_i 's are continuous, we have the following inequality:

$$|\tilde{A}(\tilde{\sigma})| \leq C \int_0^{\tilde{\sigma}} |\tilde{A}(r)| dr \quad \text{for all } \tilde{\sigma} \in [0, T/2].$$

Now by Gronwall's inequality, we have $\tilde{A}(\sigma) = 0$ for all $\tilde{\sigma} \in [0, T/2]$, which gives us $q_1(x) = q_2(x)$ for all $x \in \mathbb{R}^3$ such that $|x| \leq T/2$. This completes the proof.

ACKNOWLEDGEMENT

The author would like to thank Prof. Venky Krishnan for useful discussions. He is supported by ISIRD project from IIT Ropar and Start-up Research Grant SRG/2021/001432 from the Science and Engineering Research Board, Government of India.

REFERENCES

- [1] T. Aktosun, A. Machuca and P. Sacks; Determining the shape of a human vocal tract from pressure measurements at the lips, *Inverse Problems*, vol. 33, (2017), 115002, 33 pages.
- [2] K.P. Bube and R. Burridge; The one-dimensional inverse problem of reflection seismology. *SIAM Rev.* 25 (1983), no. 4, 497–559.
- [3] R. Burridge; The Gel'fand-Levitan, the Marchenko, and the Gopinath-Sondhi integral equations of inverse scattering theory, regarded in the context of inverse impulse-response problems. *Wave Motion* 2 (1980), no. 4, 305–323.
- [4] E. Blåsten; Well-posedness of the Goursat problem and stability for point source inverse backscattering, 2017 *Inverse Problems* 33 125003.
- [5] F.G. Friedlander; *The wave equation on a curved space-time*. Cambridge University Press, Cambridge-New York-Melbourne, 1975. Cambridge Monographs on Mathematical Physics, No. 2.
- [6] F.G. Friedlander and M. Joshi; *The Theory of Distributions*, 2nd Edition, Cambridge University Press, 1998.
- [7] I.M. Gel'fand and G. E. Shilov; *Generalized functions*. Vol. 1. Properties and operations. Translated from the 1958 Russian original by Eugene Saletan.
- [8] M. M. Lavrent'ev, V. G. Romanov and S. P. Shishat'skiĭ; *Ill-posed problems of mathematical physics and analysis*. Translated from the Russian by J. R. Schulenberger.
- [9] W. Ning and M. Yamamoto; The Gelfand-Levitan theory for one-dimensional hyperbolic systems with impulsive inputs, *Inverse Problems* 24 (2008) 025004 (19pp).
- [10] Rakesh; An inverse impedance transmission problem for the wave equation. *Comm. Partial Differential Equations* 18 (1993), no. 3-4, 583–600.

- [11] Rakesh and P. Sacks; Impedance inversion from transmission data for the wave equation. *Wave Motion* 24 (1996), no. 3, 263–274.
- [12] Rakesh; Inversion of spherically symmetric potentials from boundary data for the wave equation. *Inverse Problems* 14 (1998), no. 4, 999–1007.
- [13] Rakesh; Characterization of transmission data for Webster’s horn equation. *Inverse Problems* 16 (2000), no. 2, L9–L24.
- [14] Rakesh; An inverse problem for a layered medium with a point source. *Problems* 19 (2003), no. 3, 497–506.
- [15] Rakesh; Inverse problems for the wave equation with a single coincident source-receiver pair. *Inverse Problems* 24 (2008), no. 1, 015012, 16 pp.
- [16] Rakesh and P. Sacks; Uniqueness for a hyperbolic inverse problem with angular control on the coefficients. *J. Inverse Ill-Posed Probl.* 19 (2011), no. 1, 107–126.
- [17] Rakesh and G. Uhlmann; The point source inverse back-scattering problem. *Analysis, complex geometry, and mathematical physics: in honor of Duong H. Phong*, 279–289.
- [18] V.G. Romanov; On the problem of determining the coefficients in the lowest order terms of a hyperbolic equation. (Russian. Russian summary) *Sibirsk. Mat. Zh.* 33 (1992), no. 3, 156–160, 220; translation in *Siberian Math. J.* 33 (1992), no. 3, 497–500.
- [19] V. G. Romanov and D. I. Glushkova; The problem of determining two coefficients of a hyperbolic equation. (Russian) *Dokl. Akad. Nauk* 390 (2003), no. 4, 452–456.
- [20] V.G. Romanov; *Integral geometry and inverse problems for hyperbolic equations*, volume 26. Springer Science and Business Media, 2013.
- [21] F. Santosa and W.W. Symes; High-frequency perturbational analysis of the surface point-source response of a layered fluid. *J. Comput. Phys.* 74 (1988), no. 2, 318–381.
- [22] M.M. Sondhi; *A survey of the vocal tract inverse problem: theory, computations and experiments*. Inverse problems of acoustic and elastic waves (Ithaca, N.Y., 1984), 1–19, SIAM, Philadelphia, PA, 1984.
- [23] P.D. Stefanov; A uniqueness result for the inverse back-scattering problem. *Inverse Problems* 6 (1990), no. 6, 1055–1064.
- [24] W. W. Symes; The seismic reflection inverse problem. *Inverse Problems* 25 (2009), no. 12, 123008, 39 pp.
- [25] M. Vashisth; An inverse problems for the wave equation with source and receiver at distinct points, *Journal of Inverse and Ill-posed Problems*, Volume 27, Issue 6, Pages 835-843.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY ROPAR, RUPNAGAR, PUNJAB - 140001, INDIA.
E-MAIL: manmohanvashisth@iitrpr.ac.in